

Supplementary Table 1. Distribution of respondents by country and WHO region (n = 358)

WHO Region	Country	n	%
AFRO (n = 13)			
	Congo	4	1.1
	Madagascar	2	0.6
	Niger	2	0.6
	South Africa	2	0.6
	Sudan	2	0.6
	Uganda	1	0.3
EMRO (n = 26)			
	Egypt	1	0.3
	Iran	1	0.3
	Jordan	6	1.7
	Morocco	1	0.3
	Pakistan	11	3.0
	Qatar	4	1.1
	Tunisia	1	0.3
	United Arab Emirates	1	0.3
EURO (n = 166)			
	Austria	3	0.8
	Belgium	5	1.4
	Bosnia and Herzegovina	1	0.3
	Croatia	7	2.0
	Estonia	3	0.8
	Finland	1	0.3
	France	11	3.1
	Georgia	2	0.6
	Germany	4	1.1
	Greece	10	2.8
	Hungary	2	1.4
	Ireland	2	0.6
	Israel	3	0.8
	Italy	21	5.9
	Kazakhstan	1	0.3
	Latvia	7	2.0
	Lithuania	2	0.6
	Luxembourg	1	0.3
	Macedonia	1	0.3
	Montenegro	1	0.3
	Netherlands	10	2.8
	Norway	3	0.8
	Poland	5	1.4
	Portugal	12	3.4
	Romania	8	2.2
	Serbia	1	0.3
	Slovakia	1	0.3
	Slovenia	8	2.2
	Spain	2	0.6
	Switzerland	10	2.8
	Türkiye	10	2.8
	Ukraine	4	1.1
	United Kingdom	4	1.1
PAHO (n = 104)			
	Argentina	4	1.1
	Bolivia	1	0.3

	Brazil	5	1.4
	Canada	4	1.1
	Chile	11	3.1
	Colombia	20	5.6
	Costa Rica	1	0.3
	Dominican Republic	4	1.1
	El Salvador	1	0.3
	Guatemala	2	0.6
	Honduras	7	2.0
	Mexico	15	4.2
	Nicaragua	1	0.3
	Panamá	1	0.3
	Paraguay	2	0.6
	Perú	7	2.0
	Puerto Rico	1	0.3
	Uruguay	2	0.6
	USA	15	4.2
SEARO (n = 16)			
	Bangladesh	5	1.4
	India	7	2.0
	Indonesia	4	1.1
WPRO (n = 33)			
	Australia	12	3.4
	China	2	0.6
	Japan	1	0.3
	Malaysia	4	1.1
	Mongolia	3	0.8
	Myanmar	1	0.3
	New Zealand	1	0.3
	Philippines	5	1.4
	South Korea	1	0.3
	Taiwan	1	0.3
	Thailand	2	0.6